



Urine Drug Testing Window of Detection

drug	window of detection*
amphetamine; methamphetamine; 3,4-methylenedioxy- methamphetamine	1-2 days
barbiturates	up to 6wk
BZDs	1-3 days; up to 6wk w/ heavy use of long-acting BZDs
buprenorphine	3-4 days
cocaine	2-4 days; 10-22 days w/ heavy use
codeine	1-2 days
fentanyl	1-2 days
heroin	1-2 days
hydrocodone	2 days

hydromorphone	1-2 days
marijuana	infrequent use of 1-3 days; chronic use of up to 30 days
methadone	2-11 days
morphine	1-2 days
oxycodone	1-1.5 days
phencyclidine (PCP)	8 days
synthetic cannabinoids	single use: 72h chronic use: >72h synthetic cathinone: variable

* Detection time may vary depending on the cutoff.

Sources:

Moeller KE, et al. Clinical Interpretation of Urine Drug Tests: What Clinicians Need to Know About Urine Drug Screens. *Mayo Clin Proc.* 2017 May;92(5):774-796.

[https://www.mayoclinicproceedings.org/article/S0025-6196\(16\)30825-4/fulltext](https://www.mayoclinicproceedings.org/article/S0025-6196(16)30825-4/fulltext)

Substance Abuse and Mental Health Services Administration. *Medications for Opioid Use Disorder*. Treatment Improvement Protocol (TIP) Series 63 Publication No. PEP21-02-01-002. Rockville, MD: Substance Abuse and Mental Health Services Administration, 2021. Accessed March 28, 2024.

<https://store.samhsa.gov/sites/default/files/pep21-02-01-002.pdf>